

PLEASE ENTER ^{up} _{10/24/01}

Amendment to the Claims

1(Original). A crystalline solvate comprising: a toluene-containing epothilone B clathrate.

2(Currently amended). The crystalline solvate according to claim 1 characterized by unit cell parameters approximately equal to the following:

Cell dimensions:	a = 11.853(1) Å
	b = 10.613(2) Å
	c = 14.328(2) Å
	Volume = 1659(1) Å ³
Space group	P2 ₁
Molecules/unit cell	[[4]] 2
Density (calculated) (g/cm ³)	1.201

wherein the crystalline solvate is at a temperature of about -33°C.

3(Original). The crystalline solvate according to claim 1 wherein said crystalline solvate is characterized by peaks in a powder x-diffraction pattern at a value of two theta (CuKα λ=1.5418Å) of about 13.4, 20.2, 22.0, and 24.9, at a temperature of 23°C .

4(Original). The crystalline solvate according to claim 3 wherein said crystalline solvate is further characterized by peaks in a powder x-ray diffraction pattern at a value of two theta (CuKα λ=1.5418 Å) of about 6.7, 8.2, 11.7, 12.7, 15.0, 15.8, 16.7, 18.5, 20.9, 21.5, 24.3, 26.3, 28.5, and 30.1, at a temperature of 23°C.

5(Original). The crystalline solvate according to claim 1, which comprises about one molecule of toluene per one molecule of the epothilone B.

6(Currently amended). The crystalline solvate according to claim 5 characterized by unit cell parameters approximately equal to the following:

Cell dimensions:	a = 11.853(1) Å
	b = 10.613(2) Å
	c = 14.328(2) Å

	Volume = 1659(1) Å ³
Space group	P2 ₁
Molecules/unit cell	[[4]] 2
Density (calculated) (g/cm ³)	1.201

wherein the crystalline solvate is at a temperature of about -33°C.

7(Original). The crystalline solvate according to claim 5 wherein said crystalline solvate is characterized by peaks in a powder x-diffraction pattern at a value of two theta (CuKα λ=1.5418Å) of about 13.4, 20.2, 22.0, and 24.9, at a temperature of 23°C .

8(Original). The crystalline solvate according to claim 7 wherein said crystalline solvate is further characterized by peaks in a powder x-ray diffraction pattern at a value of two theta (CuKα λ=1.5418 Å) of about 6.7, 8.2, 11.7, 12.7, 15.0, 15.8, 16.7, 18.5, 20.9, 21.5, 24.3, 26.3, 28.5, and 30.1, at a temperature of 23°C.

9(Original). The crystalline solvate according to claim 1 characterized by: fractional atomic coordinates substantially as listed in Table 5.